



Genetic determinants associated with cfxA-positive clinical *Capnocytophaga* isolates

Zohreh Tamanai-Shacoori, Anais Dupont, Manon Auffret, Vincent Peton, Frédérique Barloy-Hubler, Elodie Ehrmann, Martine Ropert, Martine Bonnaure-Mallet, Anne Jolivet-Gougeon

► To cite this version:

Zohreh Tamanai-Shacoori, Anais Dupont, Manon Auffret, Vincent Peton, Frédérique Barloy-Hubler, et al.. Genetic determinants associated with cfxA-positive clinical *Capnocytophaga* isolates. International Journal of Antimicrobial Agents, 2015, 46 (3), pp.356-358. 10.1016/j.ijantimicag.2015.05.006 . hal-01166301

HAL Id: hal-01166301

<https://hal-univ-rennes1.archives-ouvertes.fr/hal-01166301>

Submitted on 19 Nov 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Genetic determinants associated with *cfxA*-positive clinical *Capnocytophaga* isolates

Zohreh Tamanai-Shacoori^a, Anais Dupont^a, Manon Auffret^a, Vincent Peton^a,
Frédérique Barloy-Hubler^b, Elodie Ehrmann^c, Martine Ropert^{d,e}, Martine Bonnaure-
Mallet^a, Anne Jolivet-Gougeon^{a,d,*}anne.gougeon@univ-rennes1.fr

^aEquipe de Microbiologie, EA 1254, Université de Rennes 1, Université Européenne
de Bretagne, 2 avenue du Professeur Léon Bernard, 35043 Rennes, France

^bPlateforme Amadeus, Rennes, France

^cPôle Odontologie–CHU Nice, Faculté d'Odontologie, Université de Nice Sophia-
Antipolis, Nice, France

^dPôle Biologie–CHU Rennes, 2 rue Henri Le Guilloux, 35033 Rennes, France

^eINSERM U991, 35000 Rennes, France

<PA>Equipe de Microbiologie, EA 1254, Université de Rennes 1, 2 avenue du
Professeur Léon Bernard, 35043 Rennes, France. Tel.: +33 2 23 23 43 05
fax: +33 2 23 23 49 13

Sir,

Capnocytophaga spp. have a role in the pathogenesis of various forms of
periodontal disease and systemic infections, particularly severe in neutropenic
cancer patients. The prevalence of β -lactam-resistant oral bacteria is increasing in
clinical isolates [1]. All of the reported β -lactam-resistant *Capnocytophaga* isolates
are β -lactamase-producers, but minimum inhibitory concentrations (MICs) for the
different β -lactams are variable [2]. The objective of the current study was therefore

to explain the variability in β -lactam MIC profiles in 31 *cfxA* gene-positive oral *Capnocytophaga* spp. clinical isolates with various antibiotypes. This study investigated: (i) the presence of other β -lactamase genes in addition to *cfxA* (*bla*_{CSP-1}, *cepA/cbIA* and *cfiA*); (ii) the expression level of *cfxA* in representative isolates with different antibiotic phenotypes; and (iii) the potential causes of *cfxA* expression variability, including mutation(s) in *cfxA* genes, location of the *cfxA* gene on a plasmid or the chromosome, and detection of the prevalence of mobile genetic determinants [*mobA*, *oriT*, *repA*, *ISCoc1* and transposons (Tn)] described as being involved in *cfxA* mobilisation and dissemination in *Capnocytophaga* spp. strains.

All 31 isolates were clearly identified as *Capnocytophaga gingivalis* ($n = 1$), *Capnocytophaga* spp. ($n = 1$), *Capnocytophaga ochracea* ($n = 2$), *Capnocytophaga granulosa* ($n = 3$), *Capnocytophaga leadbetteri* ($n = 3$), *Capnocytophaga* AHN9576/AHN9798/AHN8471/ChDc/ ChDCOS43 ($n = 4$) and *Capnocytophaga sputigena* ($n = 17$) by 16S rRNA gene sequencing. MIC₉₀ and MIC₅₀ values (MICs that inhibit 50% and 90% of the isolates, respectively) were all >256 mg/L for amoxicillin and first-and second-generation cephalosporins but were variable for third-generation cephalosporins. This variation in MICs for β -lactams was not due to the concomitant presence of other resistance genes: the *cepA/cbIA* and *cfiA* genes were never detected, and the *bla*_{CSP-1} gene [3] was amplified in 11/31 (35%) of *cfxA*-positive *Capnocytophaga* isolates (Fig. 1). The presence of *bla*_{CSP-1} was not significantly associated with higher MICs of cefotaxime [MIC > 16 mg/L according to Clinical and Laboratory Standards Institute (CLSI) breakpoints (<http://clsi.org/>)] compared with the presence of *cfxA* only ($P > 0.1$). In four isolates, MICs of β -lactams were low (range, <0.016–2 mg/L) with a negative nitrocefin test, despite a

positive *cfxA* PCR. In 29/31 isolates, the presence of 966 bp corresponding to the complete sequence of *cfxA* was detected (27 isolates were β -lactam-resistant but 2 were β -lactam-susceptible). PCR assay, sequencing and in silico analysis showed that the CfxA COOH-terminal region (C-ter) in two susceptible isolates was replaced by a glycosyltransferase (96% homology) for one and with a partial hypothetical efflux pump (98% homology) for the other, with 16-bp and 82-bp overlapping gene sequences, respectively. Replacement of the whole C-ter region of *cfxA* was linked to β -lactamase gene inactivation, despite a positive *cfxA* PCR in the 5' region. Of note, the C-ter region of *cfxA* appeared to be a preferentially targeted area or 'hotspot' for the acquisition of foreign genetic material.

Among the clinical isolates, 52% harboured plasmids of 3.5, 5 and/or 9 kb. PCR amplified Tn (77.4%), *ISCoc1* (61.3%), *repA* (54.8%) and *mobA* (74.2%) that was related to plasmid detection ($P < 0.05$). This was not the case for the *oriT* gene (16.1%) ($P > 0.1$) (Supplementary Table S1). In the *cfxA*-positive *Capnocytophaga* isolates, *mobA* and *repA* genes were mainly detected (100% and 94%, respectively, among plasmid-positive isolates) and related to plasmid detection ($P < 0.05$). In 74% of *mobA*-positive isolates, the *mobA* and *cfxA* region were linked by a 96-bp intergenic sequence mainly found in plasmid-positive strains ($P < 0.01$). Higher MICs of cefotaxime (MIC > 4 mg/L) [Comité de l'Antibiogramme de la Société Française de Microbiologie (CA-SFM) 2013; http://www.sfm-microbiologie.org/page/page/showpage/page_id/90.html] were significantly related to the presence of *mobA* ($P = 0.0002$), *mobA*-*cfxA* junction ($P = 0.0015$), *repA* ($P = 0.0003$) and at least one plasmid in bacterial strains ($P < 0.0001$). The presence of *bla*_{CSP-1}, *oriT*, *ISCoc1* or Tn did not significantly influence the MICs of cefotaxime.

analysis; and Adina Pascu for editorial assistance. This work was supported by the Conseil régional de Bretagne (France).

Funding: None.

Competing interests: None declared.

Ethical approval: Not required.

References

- [1] Ehrmann E, Handal T, Tamanai-Shacoori Z, Bonnaure-Mallet M, Fosse T. High prevalence of β -lactam and macrolide resistance genes in human oral *Capnocytophaga* species. J Antimicrob Chemother 2014;69:381–4.
- [2] Jolivet-Gougeon A, Tamanai-Shacoori Z, Desbordes L, Burggraeve N, Cormier M, Bonnaure-Mallet M. Genetic analysis of an Ambler class A extended-spectrum β -lactamase from *Capnocytophaga ochracea*. J Clin Microbiol 2004;42:888–90.
- [3] Guillon H, Eb F, Mammeri H. Characterization of CSP-1, a novel extended-spectrum β -lactamase produced by a clinical isolate of *Capnocytophaga sputigena*. Antimicrob Agents Chemother 2010;54:2231–4.
- [4] García N, Gutiérrez G, Lorenzo M, García JE, Píriz S, Quesada A. Genetic determinants for *cfxA* expression in *Bacteroides* strains isolated from human infections. J Antimicrob Chemother 2008;62:942–7.

- [5] Rogers MB, Bennett TK, Payne CM, Smith CJ. Insertional activation of *cepA* leads to high-level β -lactamase expression in *Bacteroides fragilis* clinical isolates. J Bacteriol 1994;176:4376–84.

Accepted Manuscript

Fig. 1. Detection of *bla*_{CSP-1}, *mobA*, *cfxA–mobA* (junction), *oriT*, *repA*, transposons (Tn), IS*Coc1* and plasmids (presence of at least one plasmid) according to different species in clinical *cfxA*-positive *Capnocytophaga* isolates.

Accepted Manuscript

FIG 1. The *bla*_{CSP-1}, *mobA*, *cfxA-mobA* (junction), *oriT*, *repA*, *Tn*, *ISCoc1* genes and plasmid detection (presence of at least one plasmid), according to different species in clinical *cfxA*-positive *Capnocytophaga* isolates.

